

Gerngross P., Götz M., Breitenmoser U., Molinari-Jobin A. & Maronde L. 2023. Categorisation of European wildcat records according to SCALP scheme. Cat News 78. 21–24. Supporting Online Material



Photo P. Gerngross



Photo P. Gerngross.



Photo P. Gerngross.



Photo KORA



Photo KORA



Photo KORA

SOM 1. Camera trap pictures showing at least two typical characteristics of a wildcat, retained as C2 record.



Photo M. Götz



Photo Progetto Lince Italia.



Photo M. Götz

SOM 1 (cont.). Camera trap pictures showing at least two typical characteristics of a wildcat, retained as C2 record.




Photo M. Götz

SOM 1 (cont.). Camera trap pictures showing at least two typical characteristics of a wildcat, retained as C2 record.



Photo KORA



Photo KORA



Photo KORA



Photo KORA



Photo KORA



Photo M. Götz

SOM 2. Camera trap pictures showing suspected hybrids, classified as C2H records. Besides at least two typical wildcat characteristics, domestic cat features are also visible.



Photo A. Weber



Photo M. Götz



Photo P. Gerngross



Photo P. Gerngross



Photo KORA

SOM 3. Pictures on which none or only one of the typical features of a wildcat are sufficiently visible, classified as C3 record. A wildcat cannot be ruled out.



Photo KORA



Photo M. Götz



Photo M. Götz



Photo M. Götz

SOM 3 (cont.). Pictures on which none or only one of the typical features of a wildcat are sufficiently visible, classified as C3 record. A wildcat cannot be ruled out.